

University of Groningen

Improving quality of maternal and perinatal care in rural Tanzania

Mooij, Robert

DOI:
[10.33612/diss.131176661](https://doi.org/10.33612/diss.131176661)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):
Mooij, R. (2020). *Improving quality of maternal and perinatal care in rural Tanzania: Safe Motherhood*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.
<https://doi.org/10.33612/diss.131176661>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



PART

III

**RESEARCH BY
DUTCH EXPATRIATE PHYSICIANS**



**THE CONTRIBUTION OF DUTCH DOCTORS
IN GLOBAL HEALTH AND TROPICAL
MEDICINE TO RESEARCH IN
GLOBAL HEALTH IN LOW- AND
MIDDLE-INCOME COUNTRIES:
AN EXPLORATION OF THE EVIDENCE**

R Mooij, EMJ Jurgens, J van Dillen and J Stekelenburg

Mooij R, Jurgens EM, van Dillen J, Stekelenburg J. The contribution of Dutch doctors in Global Health and Tropical Medicine to research in global health in low- and middle-income countries: an exploration of the evidence. Tropical doctor 2020;50:43-9.

ABSTRACT

Most medical research is conducted in high-income countries and results may not apply to low- and middle-income countries. Some expatriate physicians combine clinical duties with research. We present global health research conducted by Dutch medical doctors in Global Health and Tropical Medicine in low- and middle-income countries and explore the value of their research.

We included all research conducted in the last 30 years by medical doctors in Global Health and Tropical Medicine in a low- and middle-income country, resulting in a PhD thesis. Articles and co-authors were found through Medline. More than half of the 18 identified PhD theses concerned maternal health and obstetrics, and the majority of the research was conducted in low-income countries, mostly in rural hospitals. Over 70 local co-authors were involved. Different aspects of these studies are discussed.

INTRODUCTION

Most medical scientific research is conducted in high-income countries (HICs).^{2,3} Evidence from these studies needs to be appropriately interpreted.⁴ Some conclusions are universally applicable, but external validity depends on the setting. This means that insights from research in HICs often need local validation elsewhere. Even though attention to conducting research in low- and middle-income countries (LMICs) is increasing, it is still not a priority.^{2,5-7} Several expatriate physicians from HICs fill some of the gaps in human resources for health in LMICs.^{8,9} In addition to their clinical, managerial and teaching responsibilities, many expatriate physicians also conduct medical scientific research, generally aiming to improve the local quality of care, and less so, to contribute to advances in medical knowledge, experience and practice. Because of the growing attention in regard to health research capacity in LMICs in recent years,^{2,5,10} including the role of local co-authors^{7,11} we will elaborate on the added value of studies of expatriate physicians.

In this paper, we studied Dutch expatriate physicians, specifically medical doctors in Global Health and Tropical medicine (MDs GHTM, Box 1), to explore the contribution of this group of experts in building an evidence base relevant to LMICs. To include a clearly defined group of research and to include only larger projects, we focused on research resulting in a PhD thesis.

METHODS

Our inclusion criteria were: PhD theses published by an MD GHTM (Box 1), conducted between 1988-2018, while clinically working in an LMIC, as defined by the World Bank Country Classification (<https://datahelpdesk.worldbank.org/knowledgebase/articles/906519>). Eligible theses were identified using repositories of Dutch universities, the archives of the NVTG (Box 1) and the archives of the Working Party International Safe Motherhood and International Reproductive health (www.safemotherhood.nl). Other working parties and sections of the NVTG were contacted at an annual conference and by contacting key-informants for further information.

To double-check whether all theses had been found, we placed an enquiry in the NVTG newsletter approaching all 800 members. The seven people that responded provided information on on-going current research and no new PhD theses were mentioned.

To find out whether PhD researches resulted in publications with local co-authors, we checked PubMed (<https://www.ncbi.nlm.nih.gov/pubmed/>). Co-authors were considered to be local researchers according to their affiliations and other contributions.

RESULTS

The work of a total number of 114 researchers was reviewed for eligibility (Figure 1).

Eighteen MDs GHTM published a thesis that met the inclusion criteria (Table 1). All theses were completely available, either in print or electronic. Most (n=10) studies were in the field of maternal health and obstetrics, some of those being part of the Safe Motherhood series of PhD theses (<https://safemotherhood.nl/publicaties-safe-motherood-serie/>).¹²⁻²⁰ Studies were

1

2

3

4

5

6

7

8

9

10

In 1907 a small group of medical doctors founded the Netherlands Society for Tropical Medicine and International Health (NVTG) originally focusing on improving health care in (former) colonies – countries now classified as LMICs (www.ntvg.org). Since the late 1960s, the NVTG has offered a training programme for MDs with ambitions to work in LMICs, with a focus on clinical practice and strengthening of health systems. The previously named ‘Tropical doctor training programme’ has evolved to the current training of two clinical terms (9-12 months) in obstetrics and gynaecology, surgery, or paediatrics; a course (three months) on Global Health and Tropical Medicine; and a clinical term (six months) in an LMIC. Around 20-30 such Dutch MDs GHTM graduate each year. Typically, these doctors work for a few years in a remote setting in an LMIC, responsible for clinical tasks, as well as teaching, supervision, and management.

Box 1. Dutch medical doctors in Global Health and Tropical Medicine and the Netherlands Society for Tropical Medicine and International Health¹




Figure 1. Flow chart selection research.

conducted in five low-income, three lower-middle-income and two upper-middle-income countries. The 18 theses resulted in 125 Medline-indexed articles, which were co-authored by >70 local colleagues.

DISCUSSION

Our search identified 18 theses successfully defended in the past 30 years by MDs GHTM in LMICs, which resulted in the approval of a doctorate. Differences depending on the setting have been identified.

Studies in LMICs are usually conducted in academic settings and dissemination of the results is limited, which results in a knowledge gap in non-academic hospitals. In LMICs, differences between small hospitals and large tertiary centres can be substantial. Additionally,

Table 1. theses

Year	Country	Title	Topic	Setting	Medline entries	Local authors
1988 ²¹	Tanzania	Maternal health care in South Western highlands Tanzania	Maternal health care	2 rural hospitals in Tanzania	6	0
1988 ²²	Tanzania	Nutrition rehabilitation in the Southwestern highlands of Tanzania: A two-way learning process	Nutrition in children	Rural district and hospital in Tanzania	3	1
1995 ¹²	Tanzania	Perinatal assessment in rural Tanzania	Perinatal outcomes	1 hospital and 5 villages in a rural area in Tanzania	9	1
1999 ²³	Zambia	Childhood malnutrition in rural Zambia	Childhood malnutrition	Rural district hospital in Zambia	5	0
2003 ¹³	Ghana	Reproductive health matters in rural Ghana	Reproductive health	Rural district., 2 district hospitals in Ghana	8	6
2003 ²⁴	Kenya	Anemia in adolescent schoolgirls in Western Kenya	Anaemia in adolescents	3 rural hospitals in Kenya	4	3
2004 ¹⁴	Zimbabwe	Vaginal birth after Caesarean section in Zimbabwe and the Netherlands	Birth after Caesarean Section	Hospitals and maternity waiting home in a rural district in Zimbabwe	4	0
2004 ¹⁵	Zambia	Health care seeking behaviour and utilisation of health services in Kalabo District, Zambia	Health-care seeking behaviour	Rural community, district hospital in Zambia	6	3
2007 ²⁵	Zambia	The role of delayed umbilical cord clamping to control infant anaemia in resource-poor settings	Umbilical cord clamping	Rural hospital in Zambia and urban hospital in Libya	8	1
2009 ¹⁶	Namibia	Obstetric audit in Namibia and the Netherlands	Obstetric audit	Semi-rural hospital in Namibia	8	0
2011 ¹⁷	Malawi and Zambia	Delay in safe motherhood	Delay in safe motherhood	Rural hospital in Zambia and district hospital in Malawi	6	1
2012 ²⁶	Thailand	Susceptibility to malaria with a focus on the postpartum period	Malaria post-partum	Rural antenatal clinics in Thailand	6	10
2012 ²⁷	Thailand	Malaria in pregnancy: ultrasound studies of fetal growth	Malaria in pregnancy	Refugee and migrant clinics on the Thai Burmese border	9	12
2012 ¹⁸	Malawi	Medical mirrors – maternal care in a Malawian district	Maternal health care	District hospital in rural Malawi	11	13
2013 ¹⁹	Malawi	Health professionals and maternal health in Malawi: mortality and morbidity at district level	Maternal health professionals	District hospital in rural Malawi	5	5

1

2

3

4

5

6

7

8

9

10

Table 1. (continued)

Year	Country	Title	Topic	Setting	Medline entries	Local authors
2013 ²⁸	Sierra Leone and Rwanda	Surgical Need & Capacity in Low and Middle Income Countries	Surgical need and capacity	Population- based surveys and hospital-based studies in Sierra Leone and Rwanda	11	9
2013 ²⁰	Tanzania	Improving maternal outcome in rural Tanzania using obstetric simulation-based training	Training in obstetrics	District hospital in Tanzania	5	1
2015 ²⁹	Kenya	Cardiovascular disease prevention in the slums of Kenya	Cardiovascular disease prevention	Population-based studies in Nairobi	11	8

rural hospitals serve a different population. This means that the clinical reality in small rural hospitals in LMICs can be very different from the evidence base and that research from these hospitals is important to create local evidence and improve clinical practice. Unfortunately, for the reasons mentioned above, these results again cannot be easily generalised. It is important that settings are comparable when results are used in a different setting.

Most MDs GHTM whose research was included in this study were posted in rural district hospitals. Hence, they were in a good position to conduct research to fill the knowledge gap in non-academic settings in LMICs. Typically, they stayed in these hospitals for a longer period (often three years), which allowed them to understand the local setting, (public) health specifics and to identify local research needs. Most doctors included in our study were supported by supervisors and funders in their country of origin. They were able to establish local research partnerships, make a locally relevant research plan and complete this. Methodology, statistics and evidence-based medicine are part of the training of Dutch MD training, and the GHTM programme contains specific course-material on qualitative research methods and research in LMICs. This is useful when working together with local doctors and non-physicians with less experience in conducting research.

In most of the included studies, local health care workers also collaborated resulting, in co-authorship. Our results show >90 local participants, many of whom had no experience in research. It is hoped that these local health workers were enthused about using scientific research to evaluate and improve their clinical practice. In this way, expatriate physicians might have contributed to sustainable medical research capacity building,^{2,30,31} acting as research mentors.³² Further research might look into this in more depth. Some of the research, mostly recent, was started by local research institutions and involved a large number of local field researchers.^{19,26-29} As well as encouraging research, involving local co-authors shows authorship parity,^{33,34} and has been advocated as a requirement for HIC researchers publishing studies conducted in LMICs.¹¹ Local ownership of the studies can be encouraged by answering locally formulated research questions, by involving staff and by direct implementation of the recommendations of the studies. This will lead to improving the quality of care in the hospitals where the studies were conducted and democratisation of science.¹⁵ Expatriate physicians with clinical experience in both settings (an HIC as well as an LMIC) provide a good starting point to conduct research, as they have easy access to study populations. Mirror-studies comparing HICs with LMICs, for example on maternal mortality^{14-16,18} prove useful for both settings. The typical outsider perspective can be helpful in audit-studies.³⁵⁻³⁷ Qualitative or mixed-methods studies are appropriate types of research when little previous research is done, and new ideas are explored.^{38,39} The expatriate physician appeared to be well suited for implementation research, studying how to implement new techniques, such as best ultrasound scanning techniques, often having worked with these techniques previously in an HIC.^{27,40} This type of research is of lower-level evidence than randomised controlled trials and meta-analyses from HICs. However, the relevance of such studies is undisputed, and the results can be immediately used in the local setting to improve quality of care. New knowledge thus generated could in many cases also easily be translated to other low-resource settings. Besides

1

2

3

4

5

6

7

8

9

10

being useful for the setting in which the research is conducted, lessons may be learned for HIC settings. This is especially the case for diseases which are rare in HICs such as malaria and measles in pregnancy,^{27,41} eclampsia and uterine rupture,^{36,42} and procedures such as symphysiotomy,²¹ which are more difficult to study in HICs. Some theses have demonstrated a direct improvement of care,³⁶ and implementation of evidence-based practices, such as audits.^{15,16} The practice of delayed cord clamping has been adopted into HIC guidelines.^{43,44} About the lasting effects in the local setting after the thesis is completed, less is known.

Challenges cited by MDs GHTM in their theses were concerned with involving local health care workers too busy with clinical duties to engage in research as well as obtaining grants for relatively small studies which may seem redundant for funders without knowledge of LMIC settings. Other problems mentioned were getting local (ethical) clearance and difficulties when expensive tests (sometimes unavailable in LMICs) were needed. Implementing results into practice is a challenge for all research, but was also mentioned.

We have shown that small-scale research in low-resource settings may give useful new insights. It is important that policymakers and funding agents realise that this type of research is important in complementing research of high level of evidence in HICs. This paper shows that some Dutch expatriate physicians extend their role in their clinical field to research and we recommend continuing stimulating research and offering research methodology as a part of their training programme.⁴⁵⁻⁴⁷

We purposefully included only research by MDs GHTM resulting in a PhD. Since we only included 18 theses, which is a selection of all research by expatriate physicians, the sample might not be representative in all aspects. However, this group is well described, and for the discussion, we doubt whether a larger sample would change our conclusions. The role of PhD researchers in global health research has been described before.⁴⁸

CONCLUSION

In the last 30 years, different types of studies in LMICs have been done by MDs GHTM, resulting in 18 PhD theses. Most of the studies are in the field of maternal health and obstetrics, and >70 local colleagues were involved as co-authors. Expatriate physicians are in a unique position to conduct scientific research in an LMIC setting, in addition to their clinical and other tasks. This is of added value to the setting where the research is done, as a way of quality improvement and by building research capacity in remote areas. The country of origin of the expatriate physician benefits, as well as the physicians themselves.

ACKNOWLEDGEMENTS

A.J. van der Meulen for the initial version of the historical overview of PhD theses.

REFERENCES

1. Hummelen M. Into the world. Arnhem, The Netherlands: Uitgeverij Boekschap; 2017.
2. Beran D, Byass P, Gbakima A, et al. Research capacity building-obligations for global health partners. *The Lancet Global health* 2017;5:e567-e8.
3. Kilama WL. The 10/90 gap in sub-Saharan Africa: resolving inequities in health research. *Acta tropica* 2009;112 Suppl 1:S8-s15.
4. Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ (Clinical research ed)* 1996;312:71-2.
5. Franzen SR, Chandler C, Lang T. Health research capacity development in low and middle income countries: reality or rhetoric? A systematic meta-narrative review of the qualitative literature. *BMJ open* 2017;7:e012332.
6. Chersich MF, Martin G. Priority gaps and promising areas in maternal health research in low- and middle-income countries: summary findings of a mapping of 2292 publications between 2000 and 2012. *Globalization and health* 2017;13:6.
7. Keiser J, Utzinger J, Tanner M, Singer BH. Representation of authors and editors from countries with different human development indexes in the leading literature on tropical medicine: survey of current evidence. *BMJ (Clinical research ed)* 2004;328:1229-32.
8. Laleman G, Kegels G, Marchal B, Van der Roost D, Bogaert I, Van Damme W. The contribution of international health volunteers to the health workforce in sub-Saharan Africa. *Human resources for health* 2007;5:19.
9. Parekh N, Sawatsky AP, Mbata I, Muula AS, Bui T. Malawian impressions of expatriate physicians: A qualitative study. *Malawi medical journal : the journal of Medical Association of Malawi* 2016;28:43-7.
10. Mahawar KK, Malviya A, Kumar G. Who publishes in leading general surgical journals? The divide between the developed and developing worlds. *Asian journal of surgery* 2006;29:140-4.
11. Kushner AL, Kyamanywa P, Adisa CA, et al. Editorial policy on co-authorship of articles from low- and middle-income countries. *World journal of surgery* 2011;35:2367-8.
12. Walraven G. Perinatal assessment in rural Tanzania. Nijmegen: Radboud University; 1995.
13. Geelhoed DW. Reproductive health matters in rural Ghana. Leiden: Leiden University; 2003.
14. Spaans WA. Vaginal birth after caesarean section in Zimbabwe and the Netherlands. Amsterdam: University of Amsterdam; 2004.
15. Stekelenburg J. Health care seeking behaviour and utilisation of health services in Kalabo district, Zambia. Amsterdam: VU University; 2004.
16. van Dillen J. Obstetric audit in Namibië and the Netherlands. Amsterdam: VU University; 2009.
17. van Lonkhuijzen LRCW. Delay in safe motherhood. Groningen: Groningen University; 2011.
18. van den Akker T. Medical mirrors – maternal care in a malawian district. Amsterdam: VU University; 2012.
19. Beltman JJ. Health professionals and maternal health in Malawi: mortality and morbidity at district level. Amsterdam: VU University; 2013.
20. Nelissen EJT. Improving maternal outcome in rural Tanzania using obstetric simulation-based training. Amsterdam: VU University; 2014.
21. van Roosmalen J. Maternal health care in South Western highlands Tanzania. Leiden: Leiden University; 1988.
22. van Roosmalen-Wiebenga MW. Nutrition rehabilitation in the Southwestern highlands of Tanzania: A two-way learning process. Leiden: Leiden University; 1988.
23. Gernaat HB. Childhood malnutrition in rural Zambia. Leiden: Leiden University; 1999.
24. Leenstra T. Anemia in adolescent schoolgirls in Western Kenya. Amsterdam: University of Amsterdam; 2003.
25. van Rheenen P. The role of delayed umbilical cord clamping to control infant anaemia in resource-poor settings. Amsterdam: University of Amsterdam; 2007.

26. Boel ME. Susceptibility to malaria with a focus on the postpartum period. Amsterdam: University of Amsterdam; 2012.
27. Rijken MJ. Malaria in pregnancy: ultrasound studies of fetal growth. Utrecht: Utrecht University; 2012.
28. Groen RS. Surgical Need & Capacity in Low and Middle Income Countries. Amsterdam: University of Amsterdam; 2013.
29. van de Vijver S. Cardiovascular disease prevention in the slums of Kenya. Amsterdam: University of Amsterdam; 2015.
30. Gonzalez-Alcaide G, Park J, Huamani C, Ramos JM. Dominance and leadership in research activities: Collaboration between countries of differing human development is reflected through authorship order and designation as corresponding authors in scientific publications. *PloS one* 2017;12:e0182513.
31. Hyder AA, Rattani A, Pratt B. Research Capacity Strengthening in Low- and Middle-Income Countries: Ethical Explorations. *The Journal of law, medicine & ethics : a journal of the American Society of Law, Medicine & Ethics* 2017;45:129-37.
32. Lescano AG, Cohen CR, Raj T, et al. Strengthening Mentoring in Low- and Middle-Income Countries to Advance Global Health Research: An Overview. *The American journal of tropical medicine and hygiene* 2019;100:3-8.
33. Rees CA, Lukolyo H, Keating EM, et al. Authorship in paediatric research conducted in low- and middle-income countries: parity or parasitism? *Tropical medicine & international health : TM & IH* 2017;22:1362-70.
34. English KM, Pourbohloul B. Increasing health policy and systems research capacity in low- and middle-income countries: results from a bibliometric analysis. *Health research policy and systems* 2017;15:64.
35. van Dillen J, Stekelenburg J, Schutte J, Walraven G, van Roosmalen J. The use of audit to identify maternal mortality in different settings: is it just a difference between the rich and the poor? *Healthc Q* 2007;10:133-8.
36. van den Akker T, Mwagomba B, Irlam J, van Roosmalen J. Using audits to reduce the incidence of uterine rupture in a Malawian district hospital. *International journal of gynaecology and obstetrics: the official organ of the International Federation of Gynaecology and Obstetrics* 2009;107:289-94.
37. Heemelaar S, Nlissen E, Mdoo P, Kidanto H, van Roosmalen J, Stekelenburg J. Criteria-based audit of caesarean section in a referral hospital in rural Tanzania. *Tropical medicine & international health : TM & IH* 2016;21:525-34.
38. Stekelenburg J, Jager BE, Kolk PR, Westen EH, van der Kwaak A, Wolffers IN. Health care seeking behaviour and utilisation of traditional healers in Kalabo, Zambia. *Health policy (Amsterdam, Netherlands)* 2005;71:67-81.
39. Groen RS, Sriram VM, Kamara TB, Kushner AL, Blok L. Individual and community perceptions of surgical care in Sierra Leone. *Tropical medicine & international health : TM & IH* 2014;19:107-16.
40. Groen RS, Leow JJ, Sadasivam V, Kushner AL. Review: indications for ultrasound use in low- and middle-income countries. *Tropical medicine & international health : TM & IH* 2011;16:1525-35.
41. Mooij R, Lugumila J, Mwashambwa M. A case of unusual , complicated adult measles during a measles outbreak in Ndala Hospital , Tanzania. *Medicus Tropicus bulletin* 2013;51:18-9.
42. Mooij R, Lugumila J, Mwashambwa MY, Mwampagatwa IH, van Dillen J, Stekelenburg J. Characteristics and outcomes of patients with eclampsia and severe pre-eclampsia in a rural hospital in Western Tanzania: a retrospective medical record study. *BMC pregnancy and childbirth* 2015;15:213.
43. van Rheenen P, de Moor L, Eschbach S, de Grooth H, Brabin B. Delayed cord clamping and haemoglobin levels in infancy: a randomised controlled trial in term babies. *Tropical medicine & international health : TM & IH* 2007;12:603-16.
44. Committee Opinion No. 684: Delayed Umbilical Cord Clamping After Birth. *Obstetrics and gynecology* 2017;129:e5-e10.
45. Bezuidenhout L, Chakauya E. Hidden concerns of sharing research data by low/middle-income country scientists. *Global bioethics = Problemi di bioetica* 2018;29:39-54.

46. Bruins B. Answers to questions of member of parliament concerning the risk of losing the tropical doctor. In: Ministry of Health WaS, ed. The Hague2018.
47. Schagen van Leeuwen LH, W. Een beschaafd land zendt dokters uit. Medisch Contact 2019;74.
48. Walker SH, Ouellette V, Ridde V. How can PhD research contribute to the global health research agenda? Canadian journal of public health = Revue canadienne de sante publique 2006;97:145-8.

1

2

3

4

5

6

7

8

9

10